


# Thickness of the Middle Miocene sandy deposits of the Nordland Group

(suggested name – Eir Formation)



**Fig. 12:** Thickness of the Middle Miocene in the northern North Sea consisting mainly of sandy deposits (after Eidvin et al. 2013b). The extent of the Hutton sand is according to Gregersen & Johannessen (2007), and the extent of the Middle Miocene mudstones is according to Rundberg & Eidvin (2005).

## References

- Eidvin, T., Riis, F. & Gjeldvik, I. T., 2013b: Middle Miocene sandy deposits of the Nordland Group, northern North Sea (Suggested called Eir Formation, extent and thickness, age from fossil and Sr isotope correlations, lithology, paleobathymetry and regional correlation). NGF Abstracts and proceedings, no. 1, 2013, pp. 30-31. Poster available from the internet:  
<http://www.ngf.no/Global/Norsk/3-Publikasjoner/Presentasjoner/NGF-Vinterkonferanse-2013/Poster-2-til-NGF-vintermotet-nett.pdf>

Gregersen, U. & Johannessen, P. N., 2007: Distribution of the Neogene Utsira Sand and Hutton Sand, and the succeeding deposits in the Viking Graben area, North Sea. Marine and Petroleum Geology 24, 591-606.

Rundberg, Y. & Eidvin, T., 2005: Controls on depositional history and architecture of the Oligocene-Miocene succession, northern North Sea Basin. In B.T.G. Wandaas et al. (eds.): Onshore-Offshore Relationships on the North Atlantic Margin. NPF Special Publication 12, 207-239.